


Compound **A**: $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.32-7.26 (m, 5H), 6.86 (s, 1H), 6.85 (d, $J = 8.0$ Hz, 1H),
 6.75 (d, $J = 8.0$ Hz, 1H), 5.95 (s, 2H), 5.82 (t, $J = 8.0$ Hz, 1H), 4.77 (d, $J = 8.0$ Hz, 2H), 4.44 (s, 2H),
 3.46 (t, $J = 8.0$ Hz, 2H), 2.83 (t, $J = 8.0$ Hz, 2H), 1.49 (s, 9H)

Gao, Y.-R.; Wang, D.-Y.; Wang, Y.-Q.
Org. Lett. **2017**, *19*, 3516–3519



Spectroscopic data of **A**

IR (thin film) ν_{max} 3269, 2938, 2124, 1736 cm^{-1} .

$^1\text{H NMR}$ (400 MHz, CDCl_3)

δ 4.20 (q, $J = 7.2$ Hz, 2H), 2.59 (m, 1H), 2.53 (d, $J = 6.4$ Hz, 1H), 2.41 (dt, $J = 18.4, 5.2$ Hz, 1H), 2.26 (s, 1H),
 2.22 (m, 1H), 2.05 (m, 2H), 1.81 (m, 1H), 1.63 (m, 1H), 1.28 (t, $J = 7.2$ Hz, 3H).

$^{13}\text{C NMR}$ (101 MHz, CDCl_3) δ 200.3, 167.5, 78.2, 71.5, 61.6, 36.4, 32.5, 31.3, 30.9, 20.1, 18.7, 14.2.

